

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/350530279>

Dynamical structures of exact soliton solutions to Burgers' equation via the bilinear approach

Article in *Partial Differential Equations in Applied Mathematics* · June 2021

CITATIONS

12

READS

85

3 authors, including:



M. Belal Hossen

Uttara University

7 PUBLICATIONS 236 CITATIONS

SEE PROFILE



Harun Or Roshid

Bangladesh University of Textiles

74 PUBLICATIONS 1,117 CITATIONS

SEE PROFILE



Dynamical structures of exact soliton solutions to Burgers' equation via the bilinear approach

M. Belal Hossen^{b,*}, Harun-Or-Roshid^a, M. Zulfikar Ali^c

^a Department of Mathematics, Pabna University of Science and Technology, Bangladesh

^b Department of Computer Science & Engineering, Uttara University, Bangladesh

^c Department of Mathematics, Rajshahi University, Rajshahi, Bangladesh

ARTICLE INFO

MSC:

35K99

35P99

35P05

Keywords:

Bilinear method

Burgers' equation

Soliton solution

Rational solutions

Rogue wave

Interaction Phenomena

ABSTRACT

The Cole–Hopf transformation is used to search for exact multiple wave solutions of Burgers' equation. Particularly, solitary wave solutions and their interaction solutions are explicitly given, and a fission phenomenon is found, which is tough to be found by other traditional methods. Different solutions are presented with different combinations of elementary functions and a necessary condition for fission is proposed. Based on the bilinear formalism and with the aid of symbolic computation, we also determine two classes of rational solutions, rogue wave solutions and rogue–kink wave solutions, using various ansätze. To exhibit dynamics, we make diverse graphical analyses for the presented solutions, which show that the solutions are reliable in both mathematical physics and engineering.

1. Introduction

Soliton equations connect affluent histories of exactly solvable systems formed in mathematics, microphysics, fluid physics, cosmology, field theory, etc. To explain some physical phenomenon further, it becomes more and more important to seek exact solutions and interactions among nonlinear wave solutions. Owing to the wide applications of soliton theory in various field of natural science such as plasma physics, mathematics, biology, nuclear physics, hydrodynamics, astrophysics and geophysics, the study on integrable models has attracted much attention of many researchers. With the development of soliton theory, many kinds of powerful methods were proposed to finding the exact solutions of nonlinear partial differential equations like inverse scattering transformation,¹ (G'/G) -expansion method,² exp-function method,^{3–5} Hirota's direct method,^{6–16} Rational function method,¹⁷ the sine–cosine method¹⁸ etc. Yet some soliton models gives completely non-elastic soliton interactions when specific conditions between the wave vectors and velocities are satisfied; at a specific time, one soliton may fission to two or more solitons (soliton fission phenomena) or two or more solitons will fusion to one soliton (soliton fusion phenomena). Actually, for numerous genuine physical models such as in macromolecule material and organic membrane,¹⁹ in even-clump DNA²⁰ and in many physical fields like plasma physics, nuclear physics and so on Refs. 21–25, people have observed the same phenomena.

Wazwaz^{26–28} investigated multiple soliton solutions such type of non-elastic phenomena. Burgers' equation and Sharma–Tasso–Olver equation are such types of model in which Wang et al.²⁹ found non-elastic soliton fission and fusion phenomena with only two dispersion relations. Alam et al.,³⁰ obtain some hyperbolic, trigonometric and rational function solutions of Burgers' equation via (G'/G) -expansion method.

The main purpose of this paper is to investigate non-elastic fission phenomenon to (1+1) - dimensional Burgers' equation using the bilinear approach and a direct test function. And obtain new solitary wave and rogue wave solutions which are addressed with some figures.

This paper is organized as. In Sections 2 and 3 two types of solutions such as multiple wave (single wave, two-soliton and three-soliton) solutions and rational polynomial function solutions of the Burgers' equation are illustrated. In Sections 4 and 5, rogue wave solution and its interaction with solitary wave are demonstrated. Finally, the conclusions are presented in Section 6.

2. Multiple wave solutions of Burgers' equation and their fission

The (1+1)-dimensional Burgers' equation^{29,30} reads as

$$u_t + H u_{xx} + uu_x = 0, \quad (2.1)$$

* Corresponding author.

E-mail addresses: belalshimul14@gmail.com, belal.cse@uttarauniversity.edu.bd (M.B. Hossen).




Fig. 1. Profile of single solitary wave solution for Eq. (2.1) by choosing suitable parameters: $H = k_1 = -1, a_0 = c_1 = 1$ (a) Perspective view of the wave Eq. (2.5) (b) Perspective view of the wave Eq. (2.6).

where H is a constant to be determined later. Eq. (2.1) is a fundamental partial differential equation occurring in various areas of applied mathematics, such as fluid mechanics, nonlinear acoustics, gas dynamics, and traffic flow.

Under the dependent variable transformation

$$u = 2H(\ln f)_x. \quad (2.2)$$

Eq. (2.1) becomes the following bilinear form

$$f f_{xt} - f_x f_t + H(f f_{xxx} - f_x f_{xx}) = 0. \quad (2.3)$$

For single soliton solution of Eq. (2.1), we first consider

$$f(x, t) = a_0 + c_1 \exp(k_1 x + w_1 t), \quad (2.4)$$

and the corresponding potential field reads $v = -u_x$.

Inserting Eq. (2.4) with Eq. (2.2) into bilinear form Eq. (2.3), we can find the value of a_0, w_1 as $a_0 = \text{const.}, w_1 = -Hk_1^2$. If we substitute the value of a_0, w_1 into Eq. (2.3) along with Eq. (2.4) then we obtain following single soliton solution:

$$u(x, t) = \frac{2Hk_1 c_1 \exp(k_1(x - Hk_1 t))}{a_0 + c_1 \exp(k_1(x - Hk_1 t))}, \quad (2.5)$$

and corresponding potential field reads as

$$v(x, t) = \frac{2Hk_1^2 c_1^2 \exp(2k_1(x - Hk_1 t))}{(a_0 + c_1 \exp(k_1(x - Hk_1 t)))^2} - \frac{2Hk_1^2 c_1 \exp(k_1(x - Hk_1 t))}{a_0 + c_1 \exp(k_1(x - Hk_1 t))}. \quad (2.6)$$

In what follows, Fig. 1 presents single solitary wave solution and its potential field of Eq. (2.5) by choosing the suitable parameters.

To achieve two soliton solutions of Eq. (2.1), we just suppose

$$f(x, t) = a_0 + c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + a_{12} c_1 c_2 \exp(\xi_1 + \xi_2), \quad (2.7)$$

where $\xi_1 = k_1 x + w_1 t, \xi_2 = k_2 x + w_2 t$ and the corresponding potential field reads $v = -u_x$.

Setting Eq. (2.7) with Eq. (2.2) into the bilinear form Eq. (2.3), we gain some polynomials which are functions of the variables x and t . Equating all the coefficients of different power of exponential to zero, we can obtain a system of algebraic equations in terms of $k_1, k_2, c_1, c_2, w_1, w_2$ and a_{12} . Solving the system of algebraic equations with the help of symbolic computation system Maple, we get some solution sets of the unknown parameters which are given below.

Set-1: $a_0 = \text{const.}, a_{12} = 0, w_1 = -Hk_1^2, w_2 = -Hk_2^2$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(k_1(x - Hk_1 t)) + k_2 c_2 \exp(k_2(x - Hk_2 t))}{a_0 + c_1 \exp(k_1(x - Hk_1 t)) + c_2 \exp(k_2(x - Hk_2 t))} \right), \quad (2.8)$$

where $a_0, c_1, c_2, k_1, k_2, H$ are arbitrary constants.

The corresponding potential field read as

$$v(x, t) = 2H \left(\frac{k_1 c_1 \exp(\eta_1) + k_2 c_2 \exp(\eta_2)}{a_0 + c_1 \exp(\eta_1) + c_2 \exp(\eta_2)} \right)^2 - 2H \left(\frac{k_1^2 c_1 \exp(\eta_1) + k_2^2 c_2 \exp(\eta_2)}{a_0 + c_1 \exp(\eta_1) + c_2 \exp(\eta_2)} \right), \quad (2.9)$$

with $\eta_1 = k_1(x - Hk_1 t)$, and $\eta_2 = k_2(x - Hk_2 t)$.

From Fig. 2, the plots of the fission phenomenon between two solitary waves with the parameters choosing as $k_1 = -1, k_2 = -1.5, H = a_0 = c_1 = c_2 = 1$, we can evidently see that, after interaction one single solitary wave fission to two solitary wave and decrease wave height with smaller potential energy. i.e., at a specific time $t = 0$. That is, when the two populations living nearer to each other after interaction between them with different characteristics and tend to be a single clannish, which is completely different from the previous nature. From careful analysis of Eq. (2.8), Eq. (2.9), it concludes that, only fission phenomena occurs for all the ranges of two arbitrary parameters k_1, k_2 . Neither elastic scattering nor fusion does exist.

Set-2: $a_0 = 0, a_{12} = \text{const.}, w_1 = -H(k_1^2 + 2k_1 k_2), w_2 = -H(k_2^2 + 2k_1 k_2)$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + a_{12}(k_1 + k_2)c_1 c_2 \exp(\xi_1 + \xi_2)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + a_{12} c_1 c_2 \exp(\xi_1 + \xi_2)} \right), \quad (2.10)$$

where $\xi_1 = k_1 x - H(k_1^2 + 2k_1 k_2)t, \xi_2 = k_2 x - H(k_2^2 + 2k_1 k_2)t$ and c_1, c_2, k_1, k_2, H are arbitrary constants and the corresponding potential field reads $v = -u_x$. This solution also has the same phenomena like solution Eq. (2.10).

Set-3: $a_0 = 0, a_{12} = 0, w_1 = H(k_2^2 - k_1^2) + w_2, w_2 = \text{const.}$, then

$$u(x, t)$$




Fig. 2. Profile of two solitary wave solution for Eq. (2.1) by choosing suitable parameters: $H = -1, k_1 = 2, k_2 = -1.5, a_0 = c_1 = c_2 = 1$ (a) Perspective view of the wave Eq. (2.8) (b) Perspective view of the wave Eq. (2.9).




Fig. 3. Profile of two solitary wave solution for Eq. (2.1) by choosing suitable parameters: $H = k_2 = -2, w_2 = 0.5, k_1 = c_1 = c_2 = 1$ (a) Perspective view of the wave Eq. (2.11) (b) Perspective view of corresponding potential field.

$$= 2H \left(\frac{k_1 c_1 \exp(k_1 x + (H(k_2^2 - k_1^2) + w_2)t) + k_2 c_2 \exp(k_2 x + w_2 t)}{c_1 \exp(k_1 x + (H(k_2^2 - k_1^2) + w_2)t) + c_2 \exp(k_2 x + w_2 t)} \right), \quad (2.11)$$

where c_1, c_2, k_1, k_2, H are arbitrary constants.

The corresponding potential field reads $v = -u_x$.

From Fig. 3 solution Eq. (2.11) is completely elastic. That is to say, the soliton amplitude, wave shape and velocity remain same after non-linear collisions. So, elastic scattering, fusion and fission phenomena do not exist for any values of the parameters.

To achieve three soliton solutions of Eq. (2.1), we just suppose

$$f(x, t) = a_0 + c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3) + a_{12} c_1 c_2 \exp(\xi_1 + \xi_2) + a_{23} c_2 c_3 \exp(\xi_2 + \xi_3) + a_{13} c_1 c_3 \exp(\xi_1 + \xi_3) + a_{123} c_1 c_2 c_3 \exp(\xi_1 + \xi_2 + \xi_3), \quad (2.12)$$

where $\xi_1 = k_1 x + w_1 t, \xi_2 = k_2 x + w_2 t, \xi_3 = k_3 x + w_3 t$ and the corresponding potential field reads $v = -u_x$.

Setting Eq. (2.12) with Eq. (2.2) into the bilinear form Eq. (2.3), we gain some polynomials which are functions of the variables x and t . Equating all the coefficients of different power of exponential

to zero, we can obtain a system of algebraic equations in terms of $k_1, k_2, k_3, w_1, w_2, w_3, c_1, c_2, c_3, a_{12}, a_{23}, a_{13}$ and a_{123} . Solving the system of algebraic equations with the aid of symbolic computation system Maple, we attain the following solutions of the unknown parameters.

Set-1: $a_0 = \text{const.}, a_{12} = a_{23} = a_{13} = a_{123} = 0, w_1 = -H k_1^2, w_2 = -H k_2^2, w_3 = -H k_3^2$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(k_1 (x - H k_1 t)) + k_2 c_2 \exp(k_2 (x - H k_2 t)) + k_3 c_3 \exp(k_3 (x - H k_3 t))}{a_0 + c_1 \exp(k_1 (x - H k_1 t)) + c_2 \exp(k_2 (x - H k_2 t)) + c_3 \exp(k_3 (x - H k_3 t))} \right), \quad (2.13)$$

where $a_0, k_1, k_2, k_3, c_1, c_2, c_3, H$ are arbitrary constants.

The corresponding potential field reads $v = -u_x$.

In Fig. 4, the plots of the fusion phenomenon of the three solitons (shock waves) with the parameters choosing as: $H = a_0 = k_1 = c_1 = c_2 = c_3 = 1, k_2 = -1.5, k_3 = 2$, it is clearly shown that three single soliton fusions present one (resonant) soliton after the interaction, i.e., for a specific time $t = 0$.




Fig. 4. Profile of three solitary wave fusion solution for Eq. (2.1) by choosing suitable parameters: $H = -1, a_0 = k_1 = c_1 = c_2 = c_3 = 1, k_2 = -1.5, k_3 = 2$ (a) Perspective view of the wave Eq. (2.13) (b) Perspective view of corresponding potential field.




Fig. 5. Profile of three solitary wave fusion solution for Eq. (2.1) by choosing suitable parameters: $H = -1, a_{12} = k_1 = c_1 = c_2 = c_3 = 1, k_2 = -1.5, k_3 = 2$ (a) Perspective view of the wave Eq. (2.14) (b) Perspective view of corresponding potential field.

Set-2: $a_{12} = \text{const.}, a_0 = a_{23} = a_{13} = a_{123} = 0, w_1 = -H(k_1^2 + 2k_1k_2), w_2 = -H(k_2^2 + 2k_1k_2), w_3 = -H(k_3^2 + 2k_1k_2)$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + k_3 c_3 \exp(\xi_3) + a_{12}(k_1 + k_2) c_1 c_2 \exp(\xi_1 + \xi_2)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3) + a_{12} c_1 c_2 \exp(\xi_1 + \xi_2)} \right), \quad (2.14)$$

where $\xi_1 = k_1 x - H(k_1^2 + 2k_1k_2)t, \xi_2 = k_2 x - H(k_2^2 + 2k_1k_2)t, \xi_3 = k_3 x - H(k_3^2 + 2k_1k_2)t$, and the corresponding potential field reads $v = -u_x$ (see Fig. 5).

Set-3: $a_{23} = \text{const.}, a_0 = a_{12} = a_{13} = a_{123} = 0, w_1 = -H(k_1^2 + 2k_2k_3), w_2 = -H(k_2^2 + 2k_2k_3), w_3 = -H(k_3^2 + 2k_2k_3)$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + k_3 c_3 \exp(\xi_3) + a_{23}(k_2 + k_3) c_2 c_3 \exp(\xi_2 + \xi_3)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3) + a_{23} c_2 c_3 \exp(\xi_2 + \xi_3)} \right), \quad (2.15)$$

where $\xi_1 = k_1 x - H(k_1^2 + 2k_2k_3)t, \xi_2 = k_2 x - H(k_2^2 + 2k_2k_3)t, \xi_3 = k_3 x - H(k_3^2 + 2k_2k_3)t$, and the corresponding potential field reads $v = -u_x$.

Set-4: $a_{13} = \text{const.}, a_0 = a_{12} = a_{23} = a_{123} = 0, w_1 = -H(k_1^2 + 2k_1k_3), w_2 = -H(k_2^2 + 2k_1k_3), w_3 = -H(k_3^2 + 2k_1k_3)$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + k_3 c_3 \exp(\xi_3) + a_{13}(k_1 + k_3) c_1 c_3 \exp(\xi_1 + \xi_3)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3) + a_{13} c_1 c_3 \exp(\xi_1 + \xi_3)} \right), \quad (2.16)$$

where $\xi_1 = k_1 x - H(k_1^2 + 2k_1k_3)t, \xi_2 = k_2 x - H(k_2^2 + 2k_1k_3)t, \xi_3 = k_3 x - H(k_3^2 + 2k_1k_3)t$, and the corresponding potential field reads $v = -u_x$.

Set-5: $a_{123} = \text{const.}, a_0 = a_{12} = a_{23} = a_{13} = 0, w_1 = -H(k_1^2 + k_1k_2 + k_1k_3 + k_2k_3), w_2 = -H(k_2^2 + k_1k_2 + k_1k_3 + k_2k_3), w_3 = -H(k_3^2 + k_1k_2 + k_1k_3 + k_2k_3)$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + k_3 c_3 \exp(\xi_3) + a_{123} \Delta c_1 c_2 c_3 \exp(\xi_1 + \xi_2 + \xi_3)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3) + a_{123} c_1 c_2 c_3 \exp(\xi_1 + \xi_2 + \xi_3)} \right), \quad (2.17)$$

where $\Delta = (k_1 + k_2 + k_3), \xi_1 = k_1 x - H(k_1^2 + k_1k_2 + k_1k_3 + k_2k_3)t, \xi_2 = k_2 x - H(k_2^2 + k_1k_2 + k_1k_3 + k_2k_3)t, \xi_3 = k_3 x - H(k_3^2 + k_1k_2 + k_1k_3 + k_2k_3)t$, and the corresponding potential field reads $v = -u_x$.




Fig. 6. Profile of rational polynomial solution for Eq. (2.1) by choosing suitable parameters: $H = c = 1$, $a = b = 2$, $d = 3$ (a) Perspective view of the wave Eq. (3.3) (b) corresponding density plot.




Fig. 7. Profile of rational polynomial solution for Eq. (2.1) by choosing suitable parameters: $H = 1$, $a = 2$, $b = -1$, $c = 3$ (a) Perspective view of the wave Eq. (3.4) (b) corresponding density plot.

Set-6: $a_{123} = a_0 = a_{12} = a_{13} = a_{23} = 0$, $w_1 = \text{const.}$, $w_2 = H(k_1^2 - k_2^2) + w_1$, $w_3 = H(k_1^2 - k_3^2) + w_1$, then

$$u(x, t) = 2H \left(\frac{k_1 c_1 \exp(\xi_1) + k_2 c_2 \exp(\xi_2) + k_3 c_3 \exp(\xi_3)}{c_1 \exp(\xi_1) + c_2 \exp(\xi_2) + c_3 \exp(\xi_3)} \right), \quad (2.18)$$

where $\xi_1 = k_1 x - w_1 t$, $\xi_2 = k_2 x + (H(k_1^2 - k_2^2) + w_1)t$, $\xi_3 = k_3 x + (H(k_1^2 - k_3^2) + w_1)t$, $k_1, k_2, k_3, c_1, c_2, c_3$ are arbitrary constants and the corresponding potential field reads $v = -u_x$.

3. Rational function solutions of Burgers' equation

By a Maple computation on

$$f = \sum_{i=0}^7 \sum_{j=0}^7 C_{ij} x^i t^j \quad (3.1)$$

we notice that such type of polynomial solution for Eq. (2.3) does not allow the degree of t greater than 1 which is analytically proved by Ma.³¹ He made a hypothesis that this is true, namely, any polynomial solution f to the bilinear equation (2.3) must have the degree of t not greater than 1. In this regard, we want to consider the trial solution for the rational polynomial is of degree as follows:

$$f = R(x)t + w(x), \quad (3.2)$$

with $R(x)$ of degree one and $w(x)$ of degree at least three.

According to the bilinear form Eq. (2.3), we have to write $R(x) = apx + qb$, $w(x) = ax^3 + bx^2 + cx + d$.

Thus using Eq. (3.2) into the bilinear equation like Eq. (2.3) and solving, we achieve a class of rational polynomial solutions of the Eq. (2.1):

$$u(x, t) = \frac{2H(-6Hat + 3ax^2 + 2bx + c)}{-(6Hax + 2bH)t + ax^3 + bx^2 + cx + d}. \quad (3.3)$$




Fig. 8. Profile of Line Rogue wave (4.5) for Eq. (2.1) by choosing suitable parameters: $H = 5$, $a_1 = 2\sqrt{-1}$, $a_3 = 1$ and $a_6 = 3$. (a) Perspective view of the wave and corresponding density plot (bottom) (b) The wave propagation pattern of the wave along the x axis.

On the other hand, if we consider $R(x)$ of degree two and $w(x)$ of degree four, then setting $f = R(x)t + w(x)$ with $R(x) = pax^2qbx + lc$, $w(x) = ax^4 + bx^3 + cx^2 + dx + k$ into the bilinear form Eq. (2.3) and solving for coefficients of different powers of $x^i t^j$ for a, p, q , and l , we get $a = 0, p = \text{const.}, q = -6H, l = -2H$, which are the same results of Eq. (3.3). Now, we can conclude that in the rational polynomial solutions $w(x)$ does not support as a polynomial of degree more than 3 (see Fig. 6).

Secondly, we consider $R(x)$ of degree two and $w(x)$ of degree two, then setting $f = R(x)t + w(x)$ with $R(x) = apx + qb$, $w(x) = ax^2 + bx + c$ into the bilinear form Eq. (2.3) and solving for coefficients of different powers of $x^i t^j$ for a, p, q , we get $a = \text{const.}, p = 0, q = -\frac{2aH}{b}$. Thus

$$u(x, t) = \frac{2H(2ax + b)}{-2Hat + ax^2 + bx + c}. \quad (3.4)$$

In what follows, Fig. 7 presents exact solution of Eq. (3.4) by choosing the suitable parameters.

4. Rogue wave solutions of Burgers' equation

To get the rogue wave solutions of Eq. (2.1), assuming $f(x, t)$ in the following new form $f = 1 + g^2 + h^2$,

$$f = 1 + g^2 + h^2 \quad (4.1)$$

$$\text{with } g(x, t) = a_1 x + a_2 t + a_3,$$

$$g(x, t) = a_1 x + a_2 t + a_3 \quad (4.2)$$

$$h(x, t) = a_4 x + a_5 t + a_6, \quad (4.3)$$

where a_i , ($1 \leq i \leq 6$) are arbitrary constants to be determined later. Setting Eq. (4.1) into bilinear form Eq. (2.3), we obtain some polynomials which are functions of the variables x and t . Equating all the coefficient of x, t and the constant term to be zero, we can obtain the set of algebraic equations for a_i , ($1 \leq i \leq 6$). Solving the system with the aid of symbolic computation system Maple, gives the following relations between the parameters a_i :

$$H = H, a_1 = a_1, a_2 = 0, a_3 = a_3, a_4 = a_1 \sqrt{-1}, a_5 = 0, a_6 = a_6. \quad (4.4)$$

Therefore, substituting Eq. (4.4) and Eq. (4.1) along with Eq. (4.2) and Eq. (4.3) into Eq. (2.3) yields the following rogue wave solution,

$$u = \frac{2H \left\{ 2(xa_1 + a_3)a_1 + 2(-xa_1 + a_6\sqrt{-1})a_1 \right\}}{1 + (xa_1 + a_3)^2 + (xa_1\sqrt{-1} + a_6)^2}, \quad (4.5)$$

where the parameters satisfy the constraints (4.4). Fig. 8 shows the sketch of rogue waves for dissimilar values H, a_1, a_3, a_6 .

5. Interaction solution between Rogue wave and solitary wave

To get the interaction phenomena between line rogue wave solution and solitary wave solutions of Eq. (2.1), assuming $f(x, t)$ as a quadratic function with exponential part, that is,

$$f = 1 + g^2 + h^2 + \lambda \exp(\gamma), \quad (5.1)$$

where g and h are defined by Eq. (4.2) and Eq. (4.3), and $\gamma(x, t) = k_1 x + w_1 t, k_1, w_1$ are the constant parameters which are determined later.

Substituting Eq. (5.1) into Eq. (2.3), with the aid of symbolic computation system Maple, we can obtain the following relations among parameters:

$$H = \lambda = a_3 = a_4 = a_6 = k_1 = \text{const.}, a_1 = a_4 \sqrt{-1}, a_2 = 0, a_5 = 0, \\ w_1 = -Hk_1^2. \quad (5.2)$$

Substituting Eq. (5.1) and Eq. (5.2) into Eq. (2.3) yields the following interaction solution

$$u = \frac{2H \left\{ 2a_4 \left(-a_4 x + a_3 \sqrt{-1} \right) + 2(xa_4 + a_6)a_4 + \lambda k_1 e^{-Htk_1^2 + xk_1} \right\}}{1 + (a_4 x \sqrt{-1} + a_3)^2 + (xa_4 + a_6)^2 + \lambda e^{-Htk_1^2 + xk_1}}. \quad (5.3)$$

In what follows, Fig. 9 presents exact solution of Eq. (5.3) by choosing the suitable parameters, which can show the interaction phenomena between solitary wave and rogue wave.




Fig. 9. Interaction of Line Rogue wave and kink solution (5.3) for Eq. (2.1) by choosing suitable parameters: $H = 3, \lambda = 1.5, a_3 = 1, a_4 = 2, a_6 = 3$ and $k_1 = 1.25$. (a) Perspective view of the wave and corresponding density plot (bottom) (b) The wave propagation pattern of the wave along the x axis.

6. Conclusion

In conclusion, the (1+1)-dimensional Burgers' equation has been investigated. We have derived solitary wave solutions and their interaction solutions Burgers' equation using the Cole–Hopf transformation. From the analysis we observed that when dispersive relation exists between the wave number and the wave speed with any one $a_{ij} = 0$ or $a_{ijk} = 0$, then fission exist in the solution.

Based on the bilinear formalism, we also determine two classes of rational solutions, rogue wave solutions and rogue–kink wave solutions to the resulting equation. We observe that rational solution is worth checking if there exists a kind of Wronskian solutions to the Burgers' equation. We also conjecture that these two types of rational solutions in Eq. (3.3) and Eq. (3.4) would contain all rational solutions to the Burgers' equation. Meanwhile, the performances of the mentioned techniques are substantially powerful and absolutely reliable to search new explicit solutions of other NPDEs.

Funding statement

This research did not receive any particular offer from funding agencies.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are real glad to the referees for their constructive comments, recommendations, valuable suggestions that tremendously improved the quality of the manuscript.

References

- Deift P, Trubowitz E. Inverse scattering on the line. *Comm Pure Appl Math.* 1979;32:121–251. <http://dx.doi.org/10.1002/cpa.3160320202>.
- Roshid HO, Alam MN, Hoque MF, Akbar MA. A new extended (G'/G) -expansion method to find exact traveling wave solutions of nonlinear evolution equations. *Math Stat.* 2013;1:162–166. <http://dx.doi.org/10.13189/ms.2013.010308>.
- Shakeel M, Mohyud-din ST, Iqbal MA. Modified extended exp-function method for a system of nonlinear partial differential equations defined by seismic sea waves. *Pramana-J Phys.* 2018;91:28. <http://dx.doi.org/10.1007/s12043-018-1601-6>.
- Roshid HO, Alam MN, Akbar MA. Traveling wave solutions for fifth order (1+1)-dimensional kaup-kupershmidt equation with the help of $\exp(-\Phi(\xi))$ -expansion method. *Walailak J Sci Tech.* 2015;12:1063–1073. <http://dx.doi.org/10.14456/vol12iss7pp>.
- Roshid HO, Rahman MA. The $\exp(-\Phi(\eta))$ -expansion method with application in the (1+1)-dimensional classical Boussinesq equations. *Results Phys.* 2014;4:150–155. <http://dx.doi.org/10.1016/j.rinp.2014.07.006>.
- Roshid HO, Ma WX. Dynamics of mixed lump-solitary waves of an extended (2+ 1)-dimensional shallow water wave model. *Phys Lett A.* 2018;382:3262–3268. <http://dx.doi.org/10.1016/j.physleta.2018.09.019>.
- Ma WX. Lump solutions to the Kadomtsev–Petviashvili equation. *Phys Lett A.* 2015;379:1975–1978. <http://dx.doi.org/10.1016/j.physleta.2015.06.061>.
- Yang JY, Ma WX. Lump solutions to the BKP equation by symbolic computation. *Int J Mod Phys B.* 2016;30:1640028. <http://dx.doi.org/10.1142/S0217979216400282>.
- Ma W.X, Zhan L. Lump solutions with higher-order rational dispersion relations. *Pramana-J Phys.* 2020;94:43. <http://dx.doi.org/10.1007/s12043-020-1918-9>.
- Zhang JB, Ma WX. Mixed lump-kink solutions to the BKP equation. *Comput Math Appl.* 2017;74:591–596. <http://dx.doi.org/10.1016/j.camwa.2017.05.010>.
- Zhang JB, Zhang DJ, Shen Q. Bilinear approach for a symmetry constraint of the modified KdV equation. *Appl Math Comput.* 2011;218:4494–4500. <http://dx.doi.org/10.1016/j.amc.2011.10.030>.
- Zhang JB, Zhang JD, Chen YD. Solving the KdV equation under bargmann constraint via bilinear approach. *Commun Theor Phys.* 2010;53:211–217. <http://dx.doi.org/10.1088/0253-6102/53/2/02>.
- Hossen MB, Roshid HO, Ali MZ. Characteristics of the solitary waves and rogue waves with interaction phenomena in a (2 + 1)-dimensional Breaking Soliton equation. *Phys Lett A.* 2018;382:1268–1274. <http://dx.doi.org/10.1016/j.physleta.2018.03.016>.
- Hossen MB, Roshid HO, Ali MZ. Multi-soliton, breathers, lumps and interaction solution to the (2+1)-dimensional asymmetric Nizhnik–Novikov–Veselov equation. *Heliyon.* 2019;5:e02548. <http://dx.doi.org/10.1016/j.heliyon.2019.e02548>.
- Ma WX, Zhang Y, Tang Y. Symbolic computation of lump solutions to a combined equation involving three types of nonlinear terms. *East Asian J Appl Math.* 2020;10:732–745. <http://dx.doi.org/10.4208/eajam.151019.110420>.
- Yang JY, Ma WX, Khalique CM. Determining lump solutions for a combined soliton equation in (2+1)-dimensions. *Eur Phys J Plus.* 2020;135:494. <http://dx.doi.org/10.1140/epjp/s13360-020-00463-z>.
- Ma WX, Lee JH. A transformed rational function method and exact solutions to the 3+1 dimensional Jimbo–Miwa equation. *Chaos Solitons Fractals.* 2009;42:1356–1363. <http://dx.doi.org/10.1016/j.chaos.2009.03.043>.
- Yan C. A simple transformation for nonlinear waves. *Phys Lett A.* 1996;224:77–84. [http://dx.doi.org/10.1016/S0375-9601\(96\)00770-0](http://dx.doi.org/10.1016/S0375-9601(96)00770-0).

19. Serkin VN, Chapela VM, Percino J, Belyaeva TL. Nonlinear tunneling of temporal and spatial optical solitons through organic thin films and polymeric waveguides. *Opt Commun*. 2001;192:237–244. [http://dx.doi.org/10.1016/S0030-4018\(01\)01216-0](http://dx.doi.org/10.1016/S0030-4018(01)01216-0).
20. Hisakado M. Breather trapping mechanism in piecewise homogeneous DNA. *Phys Lett A*. 1997;227:87–93. [http://dx.doi.org/10.1016/S0375-9601\(97\)00023-6](http://dx.doi.org/10.1016/S0375-9601(97)00023-6).
21. Khater MA, Kumar D. New exact solutions for the time fractional coupled Boussinesq–Burger equation and approximate long water wave equation in shallow water. *J Ocean Eng Sci*. 2017;2:223–228. <http://dx.doi.org/10.1016/j.joes.2017.07.001>.
22. Kivshar YS, Malomed BA. Dynamics of solitons in nearly integrable systems. *Rev Modern Phys*. 1989;61:763. <http://dx.doi.org/10.1103/RevModPhys.61.763>.
23. Hossen MB, Roshid HO, Ali MZ. Modified double sub-equation method for finding complexiton solutions to the (1 + 1) dimensional nonlinear evolution equations. *Int J Appl Comput Math*. 2017;3:1–19. <http://dx.doi.org/10.1007/s40819-017-0377-6>.
24. Ono H, Nakata I. Reflection and transmission of an ion-acoustic soliton at a step-like inhomogeneity. *J Phys Soc Japan*. 1994;63:40–46. <http://dx.doi.org/10.1143/JPSJ.63.40>.
25. Ullah MS, Roshid HO, Ali MZ, Rahman Z. Dynamical structures of multi-soliton solutions to the Bogoyavlenskii's breaking soliton equations. *Eur Phys J Plus*. 2020;135(3):282. <http://dx.doi.org/10.1140/epjp/s13360-020-00289-9>.
26. Wazwaz AM. Multiple-soliton solutions for the Boussinesq equation. *Appl Math Comput*. 2007;192:479–486. <http://dx.doi.org/10.1016/j.amc.2007.03.023>.
27. Wazwaz AM. Multiple soliton solutions and rational solutions for the (2+1)-dimensional dispersive long water–wave system. *Ocean Eng*. 2013;60:95–98. <http://dx.doi.org/10.1016/j.oceaneng.2012.12.049>.
28. Wazwaz AM. Multiple soliton solutions for the integrable couplings of the KdV and the KP equations. *Cent Eur J Phys*. 2013;11:291–295. <http://dx.doi.org/10.2478/s11534-013-0183-7>.
29. Wang S, Tang X, Lou SY. Soliton fission and fusion: Burgers equation and Sharma–Tasso–Olver equation. *Chaos Solitons Fractals*. 2004;21:231–239. <http://dx.doi.org/10.1016/j.chaos.2003.10.014>.
30. Alam MN, Akbar MA, Roshid HO. Study of nonlinear evolution equations to construct traveling wave solutions via the new approach of the generalized (G'/G)-expansion method. *Math Stat*. 2013;1:102–112. <http://dx.doi.org/10.13189/ms.2013.010302>.
31. Zhang Y, Ma WX. Rational solutions to a KdV-like equation. *Appl Math Comput*. 2015;256:252–256. <http://dx.doi.org/10.1016/j.amc.2015.01.027>.